



Bazargan Kala

Fertilizer and Pesticide Application Guideline For Wheat











NUTRITECH

021-42554 | 88603949

WWW.BAZARGANKALA.COM



Time of Application 

	 Method of Application	 Application Rate	 1 Pre-Planting	 2 Tillering	 3 Stem Elongation	 4 Heading	 5 Grain Filling
 Soil Booster 510	Irrigation	10 Lit/ha					
 Amino Chelate Fe 510	Foliar	1-2 lit/1000 L					
 K60	Foliar	1-2 lit/1000 L					
 Amino Chelate Manganese 510	Foliar	1-2 lit/1000 L					
 MicroMix 222	Foliar	1-2 lit/1000 L					
 Salt Gone 1303	Irrigation	10-20 Lit/ha					
 Amino Chelate Zn 710	Foliar	1-2 lit/1000 L					
 AminoPotasse 1010	Foliar	2-3 lit/1000 L					
 Grain Set 615 *	Foliar	5-6 lit/1000 L					
 Humic Acid 123	Seed treatment	1-1.5 lit per 100 Kg of seeds					
 Humic Acid 123	Irrigation	10 Lit/ha					
 Nitro Micro 2505	Irrigation	10-20 Lit/ha					
 High K 5-10-40	Foliar	2-3 lit/1000 L					
 High P 10-40-10	Foliar	2-3 lit/1000 L					
 Balance 20-20-20	Foliar	2-3 lit/1000 L					
 Granule 486	Soil Incorporation	200-400 kg/ha					
 Growth Mixture 1011 **	Seed treatment	1% solution					
 Growth Mixture 1011	Foliar	1-3 lit/1000 L					

* If needed to mixed with Zn, use the rate of 1.5/1000 of this product | ** Amount of water depends on the size and quantity of the seeds



Cautions & Disclaimers:

If needed to be mixed with any other products, jar test and small- scale application are highly recommended. | The dosage and recommendations above are general; In order to be exact and most effective, soil and tissue analysis should be considered and also consultation of local experienced agronomists is highly recommended. | The liability of the producer is limited to the guaranteed content and physical characteristics as claimed; the producer shall not be liable for anything out of its control. | This product is NOT miscible with alkaline solutions, sulfur products highly acidic and pesticides and shall not be mixed with the mentioned products. | The product shall be kept in 5-30 degrees centigrade and also NOT under the direct sunlight and away from flammable materials. | Foliar application shall be applied in the coolest time of the day during the day time (early morning or near sunset). | Use acceptable quality of water for foliar applications and fertigation.



Bazargan Kala

021-42554 | 88603949

WWW.BAZARGANKALA.COM

NUTRITECH®



Time of Application

	 Method of Application	 Application Rate	 1 Pre-Planting	 2 Tillering	 3 Stem Elongation	 4 Heading	 5 Grain Filling
Urea Phosphate 17-44	Irrigation	5-20 Lit/ha					
Urea Phosphate 17-44	Foliar	2-3 lit/1000 L					
Potassium Phosphite 2724	Irrigation	5-7 Lit/ha					
Potassium Phosphite 2724	Foliar	2-3 lit/1000 L					
Seed Treatment 1010 *	Seed Treatment	3-4 lit per 1 metric ton of Seeds					
pH Reduce 401 **							
Surfactant 301 ***							

* Dilute each liter of this product in maximum 2 to 5 liters of water, then mix it with the seeds until they are uniformly covered. When seeds are dried under shade, they are ready to be planted. | ** Irrigation 3-5 Lit/ha; Descaling of irrigation system 3-5 Lit/ha; Tank mix for spraying (with fertilizers and pesticides) 0.5-1 Lit/ha | *** 1/1000 Mix with pesticides and foliar fertilizers | 2/1000 Washing the plant and reducing pests' population



Cautions & Disclaimers:

If needed to be mixed with any other products, jar test and small- scale application are highly recommended. | The dosage and recommendations above are general; In order to be exact and most effective, soil and tissue analysis should be considered and also consultation of local experienced agronomists is highly recommended. | The liability of the producer is limited to the guaranteed content and physical characteristics as claimed; the producer shall not be liable for anything out of its control. | This product is NOT miscible with alkaline solutions, sulfur products highly acidic and pesticides and shall not be mixed with the mentioned products. | The product shall be kept in 5-30 degrees centigrade and also NOT under the direct sunlight and away from flammable materials. | Foliar application shall be applied in the coolest time of the day during the day time (early morning or near sunset). | Use acceptable quality of water for foliar applications and fertigation.